

POSITIVITY OF EQUIVARIANT SCHUBERT CLASSES THROUGH MOMENT MAP DEGENERATION

CATALIN ZARA

ABSTRACT. For a flag manifold $M = G/B$ with the canonical torus action, the T -equivariant cohomology is generated by equivariant Schubert classes, with one class τ_u for every element u of the Weyl group W . These classes are determined by their restrictions to the fixed point set $M^T \simeq W$, and the restrictions are polynomials with nonnegative integer coefficients in the simple roots.

The main result of this article is a positive formula for computing $\tau_u(v)$ in types A, B, and C. To obtain this formula we identify G/B with a generic co-adjoint orbit and use a result of Goldin and Tolman to compute $\tau_u(v)$ in terms of the induced moment map. Our formula, given as a sum of contributions of certain maximal ascending chains from u to v , follows from a systematic degeneration of the moment map, corresponding to degenerating the co-adjoint orbit. In type A we prove that our formula is manifestly equivalent to the formula announced by Billey in [Bi], but in type C, the two formulas are not equivalent.

CONTENTS

1. Introduction	1
2. Combinatorial Schubert Classes	3
3. Positive Formula Using Subwords	5
4. The Goldin-Tolman Formula	6
5. Limits and Chain Contributions	9
6. Positive Formula Using Chains	11
7. Type A Positive Formula Using Chains	14
8. Chains and Subwords	16
9. Equivalence of Positive Formulas	18
References	20

1. INTRODUCTION

Let G be a connected, complex, semisimple Lie group, $B \subset G$ a Borel subgroup, corresponding to a set of simple roots $\{\alpha_1, \dots, \alpha_n\}$. Let $T^{\mathbb{C}} \subset B$

be a maximal torus, and $T \subset T^{\mathbb{C}}$ a compact real form of $T^{\mathbb{C}}$. The torus T acts on the flag manifold $M = G/B$ by left multiplication on G , and the fixed point set M^T corresponds bijectively with the Weyl group W .

The equivariant cohomology ring $H_T^*(M) = H_T^*(M, \mathbb{Q})$ is a free module over $H_T^*(pt, \mathbb{Q}) \simeq \mathbb{Q}[\alpha_1, \dots, \alpha_n]$, and that is also true for integral, and not just rational coefficients. A basis of this module is given by equivariant Schubert classes, $\{\tau_u\}_{u \in W}$, with a class τ_u for each fixed point $u \in W = M^T$. The equivariant Schubert class $\tau_u \in H_T^*(M)$ is the class induced by the T -equivariant cycle $X_u = \overline{B_- u B}/B$, where B_- is the opposite Borel subgroup. The pull-back map $H_T^*(M) \rightarrow H_T^*(M^T)$ is injective, so an equivariant class is determined by its values on M^T .

In particular, τ_u is determined by the values $\tau_u(v) \in \mathbb{Q}[\alpha_1, \dots, \alpha_n]$, for $v \in W$. The integral positivity property of the equivariant Schubert class τ_u states that for all $v \in W$ we have

$$\tau_u(v) \in \mathbb{Z}_{\geq 0}[\alpha_1, \dots, \alpha_n].$$

In other words, $\tau_u(v)$ can be written as a polynomial with *nonnegative integer* coefficients in the simple roots $\alpha_1, \dots, \alpha_n$. Billey [Bi] gives an *integral positive formula* for $\tau_u(v)$: a formula for $\tau_u(v)$ as a sum of polynomials in $\alpha_1, \dots, \alpha_n$ with non-negative integer coefficients.

The main result of this article, formulated in Theorem 6.1, is a new positive formula for computing $\tau_u(v)$ in types A , B , and C . To obtain this formula we identify G/B with a generic co-adjoint orbit and use a recent result of Goldin and Tolman ([GT]) to compute $\tau_u(v)$ in terms of the induced moment map. Our formula, given as a sum of contributions of certain maximal ascending chains from u to v , follows from a systematic degeneration of the moment map, corresponding to degenerating the co-adjoint orbit. The resulting formula is integral in types A and C , but only rational in type B (and for G_2). In type A our formula is an alternative formulation of the positive announced by Billey, but in type C the two formulas, while giving the same answer, are not manifestly equivalent.

The main methods in this paper involve the combinatorics of Weyl groups. The type A formula has been circulated earlier, and motivated a separate project by Sabatini and Tolman ([ST]). Their geometrical methods allowed them to independently obtain, as applications, the same formulas for types B and C , but from a different perspective.

The content of the paper is organized as follows. In Section 2 we give a combinatorial description of τ_u and in Section 3 we recall Billey's positive formula, that uses subwords of a reduced word for v . In Section 4 we apply the result of Goldin and Tolman to co-adjoint orbits and obtain a formula for $\tau_u(v)$ in terms of maximal ascending chains. This formula depends on the particular generic co-adjoint orbit and in Section 5 we use moment

map/orbit degeneration to get a simpler formula, in which only some chains have non-zero contributions. In Section 6 we prove that the simpler formula is a positive formula, integral in types A and C and rational in type B , and in Section 7 we give an explicit version in type A . We prove the equivalence of the chain formula with the subword formula in Section 9, after first constructing, in Section 8, a general map connecting chains and subwords.

I would like to thank Rebecca Goldin, Sue Tolman, Silvia Sabatini, and Allen Knutson for explaining their work, and to all of the above and Victor Guillemin for many stimulating discussions and valuable comments while writing this paper. Many thanks are also due to John Stembridge for writing the *Maple* packages `coxeter` and `weyl`, which have been used to verify several conjectured formulas before proving them.

2. COMBINATORIAL SCHUBERT CLASSES

The injective morphism $H_T^*(M) \rightarrow H_T^*(M^T)$ identifies $H_T^*(M)$ with a subring of $H_T^*(M^T) = \text{Maps}(W, \mathbb{S})$, where $\mathbb{S} = \mathbb{Q}[\alpha_1, \dots, \alpha_n]$. Not all maps represent classes in $H_T^*(M)$, and the ones that do can be described using a discrete structure involving a regular graph $\Gamma = (V, E)$, and a labeling of the oriented edge of Γ by elements of $\mathfrak{t}^*$, the dual of the Lie algebra of T .

The vertices V of Γ correspond to the Weyl group W . Two vertices u and v are joined by an edge if and only if they differ by a reflection s_β , for some positive root β . Note that $us_\beta = s_{u\beta}u$, so if u and v differ by a reflection to the left, they also differ by a reflection to the right. If $v = us_\beta$, with $\beta \succ 0$ a positive root, then the oriented edge $e = (u, v)$ of Γ is labeled by $\alpha(u, v) = u\beta$ and is called *ascending* if $\alpha(u, v) \succ 0$. We will also use $\alpha(e)$, α_e , or $\alpha_{u,v}$ for $\alpha(u, v)$. The pair (Γ, α) is called the *GKM graph* of (M, T) .

An assignment $f: W \rightarrow \mathbb{S} = \mathbb{Q}[\alpha_1, \dots, \alpha_n]$ is a *cohomology class* if

$$f(v) - f(u) \in \alpha(e)\mathbb{S},$$

for every edge $e = [u, v]$ of Γ . The class f is called homogeneous of degree k if, for every $u \in W$, the polynomial $f(u)$ is homogeneous of degree k . The *cohomology ring* $H_\alpha^*(\Gamma)$ is the graded subring of $\text{Maps}(W, \mathbb{S})$ consisting of all classes. Then

$$H_T^{2k}(M) \simeq H_\alpha^k(\Gamma),$$

and since $H_T^{\text{odd}}(M) = 0$, that means that $H_T^*(M) \simeq H_\alpha^*(\Gamma)$. We will in general identify a class $f \in H_T^*(M)$ with its image in $H_\alpha^*(\Gamma)$. In particular, we will denote by τ_u both the equivariant Schubert class in $H_T^*(M)$ and its image in $H_T^*(\Gamma)$. Sometimes we will refer to $\tau_u \in H_T^*(\Gamma)$ as the *combinatorial Schubert class*.

To give the combinatorial description of the (combinatorial) Schubert class $\tau_u \in H_\alpha^*(\Gamma)$, we start by recalling some results concerning the combinatorics

of the Weyl group W . Most results are valid for general Coxeter groups, and details can be found, for example, in [BB].

Let R be a root system of rank n , let W be its Weyl group, and let $\mathcal{B} = \{\alpha_1, \dots, \alpha_n\}$ be a choice of simple roots. Then $\mathcal{B}$ is a basis of $\mathfrak{t}^*$, the dual of the Lie algebra of T . For a non-zero vector $\beta \in \mathfrak{t}^*$ we say that $\beta \succ 0$ if β is in the non-negative cone over $\{\alpha_1, \dots, \alpha_n\}$, in other words, if the coordinates of β in the basis $\mathcal{B}$ are non-negative.

For $i = 1, \dots, n$, let $s_i = s_{\alpha_i} \in W$ be the reflection generated α_i . A word of length m is an array $I = [i_1, i_2, \dots, i_m]$ with entries (letters) from $\{0, 1, \dots, n\}$. To each nonempty word I we associate the Weyl group element $s_I = s_{i_1} s_{i_2} \dots s_{i_m}$. (If the word I is empty, then s_I is the identity.) A subword of a word I is a word $J = [\epsilon_1 i_1, \epsilon_2 i_2, \dots, \epsilon_m i_m]$, with $\epsilon_k \in \{0, 1\}$ for all $k = 1, \dots, m$. A word I is *reduced* if $s_J \neq s_I$ for all subwords J of I other than I itself. If $v \in W$, then $v = s_{i_1} s_{i_2} \dots s_{i_m}$ is called a *decomposition* for v , and $I = [i_1, i_2, \dots, i_m]$ is a word for v . The decomposition is *reduced* if the word is reduced. All reduced words for $v \in W$ have the same number of letters, and this common number, denoted by $\ell(v)$, is the *length* of v .

We define a partial order on W as follows. If $v = us_\beta$, with $\beta \succ 0$, we define $u \prec v$ if $u\beta \succ 0$. The (*strong*) *Bruhat order* on W is the transitive closure of $\prec$. In other words, $u \preceq v$ if and only if there exists an ascending chain

$$(1) \quad u = u_0 \xrightarrow{*s_{\beta_1}} u_1 \xrightarrow{*s_{\beta_2}} u_2 \longrightarrow \dots \xrightarrow{*s_{\beta_m}} u_m = v .$$

where $p \xrightarrow{*s_\beta} q$ means $q = ps_\beta$. An equivalent definition can be given in terms of words and subwords: $u \preceq v$ if and only if every reduced word I for v has a subword that is a word for u . Let $\mathcal{F}_u = \{v \mid u \preceq v\}$ be the flow-up from u under the strong Bruhat order.

For $u \in W$, we define

$$\Lambda_u^- = \prod \{\beta \mid \beta \succ 0 \text{ and } u^{-1}\beta \prec 0\}$$

If $I = [i_1, \dots, i_m]$ is a reduced word for v , then the positive roots that are sent by v^{-1} into negative roots are

$$\alpha_{i_1} , \quad s_{i_1} \alpha_{i_2} , \quad s_{i_1} s_{i_2} \alpha_{i_3} , \quad \dots , \quad s_{i_1} \dots s_{i_{m-1}} \alpha_{i_m} ,$$

hence

$$\Lambda_v^- = \prod_{j=1}^m s_{i_1} s_{i_2} \dots s_{i_{j-1}} \alpha_{i_j} .$$

Example 2.1. We illustrate the results above for $G = SL_n(\mathbb{C})$, the special linear group of complex matrices with determinant equal to one, with the Borel subgroup B being the subgroup of upper triangular matrices. Then

$M = G/B$ is the manifold of complete flags in $\mathbb{C}^n$. The fixed points of the T -action are the classes of permutation matrices, hence M^T is in a bijective correspondence with $W = S_n$, the set of permutations of $[n] = \{1, \dots, n\}$. We use the one-line notation $u = u(1)u(2)\dots u(n)$ for permutations. The simple roots are $\alpha_1 = x_1 - x_2, \dots, \alpha_{n-1} = x_{n-1} - x_n$, the positive roots are $\alpha_{ij} = x_i - x_j = \alpha_i + \dots + \alpha_{j-1}$, for $1 \leq i < j \leq n$, and the Weyl group $W = S_n$ acts on roots by permuting the x variables. The reflection $s_{\alpha_{ij}}$ corresponds to the transposition (i, j) that swaps i and j ; if $j = i+1$, we denote $s_{\alpha_{ij}}$ by s_i and call it a *simple* transposition. By convention, s_0 is the identity of S_n .

For a permutation $v \in S_n$, the set of positive roots that are sent by v^{-1} into negative roots corresponds bijectively to the set of inversions-as-values,

$$\text{Inv}(v) = \{(v(j), v(i)) \mid i < j, v(i) > v(j)\}.$$

If $I = [i_1, \dots, i_m]$ is a reduced word for v , then

$$\Lambda_v^- = \prod_{(a,b) \in \text{Inv}(v)} (x_a - x_b) = \prod_{k=1}^m s_{i_1} \cdots s_{i_{k-1}} \alpha_{i_k}.$$

If $v = u(i, j)$ with $i < j$, then $u \prec v$ in the strong Bruhat order if and only if $u(i) < u(j)$. In other words, $u \prec v = u(i, j)$ if $(u(i), u(j)) \notin \text{Inv}(u)$, or, equivalently, if $(v(i), v(j)) \in \text{Inv}(v)$.

The description of the combinatorial Schubert class τ_u is the following.

Proposition 2.2. For every $u \in W$, the class $\tau_u \in H_\alpha^*(\Gamma)$ is the unique class satisfying the following properties:

- (1) τ_u is homogeneous, of degree $\ell(u)$, the length of u ;
- (2) τ_u is supported on $\mathcal{F}(u)$, the flow-up from u ;
- (3) τ_u is normalized by $\tau_u(u) = \Lambda_u^-$.

3. POSITIVE FORMULA USING SUBWORDS

In this section we recall Billey's integral positive formula. Let $u, v \in W$, with $u \preceq v$, and let $I = [i_1, i_2, \dots, i_m]$ be a reduced word for v . For a subword $J = [\epsilon_1 i_1, \epsilon_2 i_2, \dots, \epsilon_m i_m]$ of I , define the *subword contribution* $SC(J, I)$ as the following product of positive roots:

$$\begin{aligned} SC(J, I) &= \prod_{j=1}^m [s_{i_1} s_{i_2} \cdots s_{i_{j-1}} \alpha_{i_j}]^{\epsilon_j} = \prod_{\substack{j=1 \\ \epsilon_j=1}}^m s_{i_1} s_{i_2} \cdots s_{i_{j-1}} \alpha_{i_j} = \\ &= \Lambda_v^- \cdot \prod_{\substack{j=1 \\ \epsilon_j=0}}^m \frac{1}{s_{i_1} s_{i_2} \cdots s_{i_{j-1}} \alpha_{i_j}}, \end{aligned}$$

hence $SC(J, I)$ is obtained from Λ_v^- by canceling the positive roots generated by the deleted letters ($\epsilon_j = 0$). Let $\mathcal{R}(u, I)$ be the set of subwords of I that, after deleting the zeroes, become *reduced* words for u .

Theorem 3.1 ([Bi]). Let $u, v \in W$ and let $I = [i_1, \dots, i_m]$ be a reduced word for v . The value at v of the Schubert class τ_u is given by

$$(2) \quad \tau_u(v) = \sum_{J \in \mathcal{R}(u, I)} SC(J, I) = \sum_{J \in \mathcal{R}(u, I)} \prod_{\substack{j=1 \\ \epsilon_j=1}}^m s_{i_1} s_{i_2} \cdots s_{i_{j-1}} \alpha_{i_j}.$$

Example 3.2. Let $u = 2143$ and $v = 3421$ in S_4 . A reduced word for v is $I = [2, 1, 3, 2, 3]$. There are two subwords of I that are reduced words for u after deleting all the zeroes: $J_1 = [0, 1, 3, 0, 0]$ and $J_2 = [0, 1, 0, 0, 3]$. Their contributions are

$$\begin{aligned} SC(J_1, I) &= [\alpha_2]^0 \cdot [s_2 \alpha_1]^1 \cdot [s_2 s_1 \alpha_3]^1 \cdot [s_2 s_1 s_3 \alpha_2]^0 \cdot [s_2 s_1 s_3 s_2 \alpha_3]^0 = \\ &= [s_2 \cdot (x_1 - x_2)] \cdot [s_2 s_1 \cdot (x_3 - x_4)] = (x_1 - x_3)(x_2 - x_4) = \\ &= (\alpha_1 + \alpha_2)(\alpha_2 + \alpha_3) \\ SC(J_2, I) &= [\alpha_2]^0 \cdot [s_2 \alpha_1]^1 \cdot [s_2 s_1 \alpha_3]^0 \cdot [s_2 s_1 s_3 \alpha_2]^0 \cdot [s_2 s_1 s_3 s_2 \alpha_3]^1 = \\ &= [s_2 \cdot (x_1 - x_2)] \cdot [s_2 s_1 s_3 s_2 \cdot (x_3 - x_4)] = (x_1 - x_3)(x_1 - x_2) = \\ &= (\alpha_1 + \alpha_2) \alpha_1. \end{aligned}$$

Therefore

$$\tau_{2143}(3421) = SC(J_1, I) + SC(J_2, I) = (\alpha_1 + \alpha_2)(\alpha_1 + \alpha_2 + \alpha_3).$$

4. THE GOLDIN-TOLMAN FORMULA

Goldin and Tolman ([GT]) have recently announced a formula for computing the values of what they call *canonical classes*, valid for more general spaces¹. For flag manifolds, their classes satisfy conditions 1 and 3 above, but condition 2 is replaced by $\tau_u(v) = 0$ if $\ell(v) \leq \ell(u)$ and $v \neq u$. However, since the length function is strictly increasing with respect to the Bruhat order (that is, if $u \prec v$ then $\ell(u) < \ell(v)$), it turns out that in this case the canonical generators are the equivariant Schubert classes, and their formula can be used to compute the values $\tau_u(v)$ for $u, v \in W$.

The Goldin-Tolman formula involves two more ingredients. The first ingredient is the subgraph Γ_0 of Γ , having the same vertices as Γ , but only the edges $e = (u, v)$ of Γ for which $\ell(v) = \ell(u) \pm 1$. If u and v are elements in W , let $\Sigma(u, v)$ be the set of ascending chains in Γ_0 from u to v . These chains are the maximal length ascending chains in Γ from u to v . Every edge of

¹Allen Knutson has informed me that the flag manifold case of this formula also appeared, in implicit form, in his earlier paper [Kn].

such a chain will be considered oriented, with the orientation that makes it an ascending edge.

The second ingredient is a moment map. The Lie algebra $\mathfrak{g}$ of G can be canonically identified with its dual $\mathfrak{g}^*$ using the Killing form, and that allows us to regard $\mathfrak{t}^*$ as a subspace of $\mathfrak{g}^*$. Let $\eta \in \mathfrak{t}^* \subset \mathfrak{g}^*$ be in the interior of the positive Weyl chamber, and let $\mathcal{O}_\eta = G \cdot \eta \subset \mathfrak{g}^*$ be the co-adjoint orbit through η . Then the stabilizer of η is B , and hence $\mathcal{O}_\eta \simeq G/B = M$ as T -spaces.

The co-adjoint orbit $\mathcal{O}_\eta$ is a Hamiltonian G -space, with moment map given by the inclusion $\mathcal{O}_\eta \hookrightarrow \mathfrak{g}^*$. Therefore it is also a Hamiltonian T -space, with moment map given by inclusion followed by projection onto $\mathfrak{t}^*$. With the symplectic structure induced by the identification $G/B \simeq \mathcal{O}_\eta$, the flag manifold $M = G/B$ is a Hamiltonian T -space, with moment map $\phi_\eta: G/B \rightarrow \mathfrak{t}^*$. If P_w is the fixed point corresponding to the element $w \in W$ of the Weyl group, then $\phi_\eta(P_w) = w \cdot \eta$. Identifying the fixed point P_w with the Weyl group element w , we get a map $\phi_\eta: W \rightarrow \mathfrak{t}^*$, given by $\phi_\eta(w) = w\eta$.

Applying the Goldin-Tolman formula ([GT]), we get the following result.

Theorem 4.1. If $\eta \in \mathfrak{t}^*$ is in the positive Weyl chamber, then

$$(3) \quad \tau_u(v) = \sum_{\gamma \in \Sigma(u,v)} E_\eta(\gamma) ,$$

where, for every ascending chain of maximal length

$$(4) \quad \gamma : u = u_0 \longrightarrow u_1 \longrightarrow \cdots \longrightarrow u_m = v$$

in $\Sigma(u, v)$, the contribution $E_\eta(\gamma)$ is given by

$$(5) \quad E_\eta(\gamma) = \Lambda_v^- \prod_{k=1}^m \left(\frac{1}{\alpha(u_{k-1}, u_k)} \cdot \frac{\phi_\eta(u_k) - \phi_\eta(u_{k-1})}{\phi_\eta(v) - \phi_\eta(u_{k-1})} \right) .$$

There are two important features in this formula. The first is that each term depends on η , but the sum doesn't. The second is that each term is a rational expression, but the sum is a polynomial. Therefore a lot of cancelations must occur when summing, and a first indication of how that happens is the following.

Lemma 4.2. If $p, q \in W$ and $q = ps_\beta$ with $\beta \succ 0$, then

$$(6) \quad \frac{\phi_\eta(q) - \phi_\eta(p)}{\alpha(p, q)} < 0 .$$

Proof.

$$\frac{\phi_\eta(q) - \phi_\eta(p)}{\alpha(p, q)} = \frac{ps_\beta\eta - p\eta}{p\beta} = -\langle \eta, \beta \rangle = -\frac{2(\eta, \beta)}{(\beta, \beta)} < 0 ,$$

since β is a positive root and η is in the positive Weyl chamber. $\square$

Hence, if γ is the maximal chain (4) and $u_k = u_{k-1}s_{\beta_k}$ with $\beta_k \succ 0$, then

$$E_\eta(\gamma) = \Lambda_v^- \prod_{k=1}^m \frac{\langle \eta, \beta_k \rangle}{u_{k-1}\eta - v\eta}.$$

Lemma 4.3. Let $p \prec q$, let η be in the closure of the positive Weyl chamber and let $\beta = p\eta - q\eta$. Then $\beta \succcurlyeq 0$. If $\beta \succ 0$, then $p^{-1}\beta \succ 0$ and $q^{-1}\beta \prec 0$.

Proof. If $p \prec q$, then there exists an ascending chain

$$p = p_0 \xrightarrow{*s_{\beta_1}} p_1 \xrightarrow{*s_{\beta_2}} p_2 \rightarrow \cdots \xrightarrow{*s_{\beta_m}} p_m = q$$

with $\beta_k \succ 0$ and $p_{k-1}\beta_k \succ 0$. Then

$$\begin{aligned} (7) \quad p\eta - q\eta &= (p\eta - p_1\eta) + \cdots + (p_{k-1}\eta - p_k\eta) + \cdots + (p_{m-1}\eta - q\eta) = \\ &= \langle \eta, \beta_1 \rangle p_0\beta_1 + \cdots + \langle \eta, \beta_k \rangle p_{k-1}\beta_k + \cdots + \langle \eta, \beta_m \rangle p_{m-1}\beta_m, \end{aligned}$$

and each of the terms in the last sum is in the non-negative cone generated by the simple roots, hence $\beta \succcurlyeq 0$. Moreover, $\beta = 0$ if and only if all the factors $\langle \eta, \beta_k \rangle$ are zero.

We have $p^{-1}\beta = \eta - p^{-1}q\eta = id \eta - p^{-1}q\eta$, where id is the identity in the Weyl group. Since $id \prec p^{-1}q$, if $\beta \succ 0$ then $p^{-1}\beta \succ 0$, and similarly, $q^{-1}\eta = -(id \eta - q^{-1}p\eta) \prec 0$. $\square$

Therefore $E_\eta(\gamma)$ is a ratio of two polynomials in the simple roots $\alpha_1, \dots, \alpha_n$, homogeneous and with non-negative coefficients. Moreover, if $u_{k-1}\eta - v\eta$ is a multiple of a root, then it cancels (over $\mathbb{R}_{>0}$) one of the factors of Λ_v^- .

Let $\{\omega_1, \dots, \omega_n\}$ be the fundamental weights corresponding to the simple roots $\{\alpha_1, \dots, \alpha_n\}$, defined by the conditions

$$\langle \omega_i, \alpha_j \rangle = \delta_{ij} \quad \text{or, equivalently,} \quad (\omega_i, \alpha_j) = \frac{1}{2}(\alpha_i, \alpha_i)\delta_{ij}.$$

Then $\{\omega_1, \dots, \omega_n\}$ is a basis of $\mathfrak{t}^*$, and $\eta = \mu_1\omega_1 + \cdots + \mu_n\omega_n \in \mathfrak{t}^*$ is in the positive Weyl chamber if and only if $\mu_j > 0$ for all $j = 1, \dots, n$.

Then the contribution of the chain γ is

$$(8) \quad E_\mu(\gamma) = E_{\mu_1\omega_1 + \cdots + \mu_n\omega_n}(\gamma) = \Lambda_v^- \prod_{k=1}^m \frac{\sum_{i=1}^n \langle \omega_i, \beta_k \rangle \mu_i}{\sum_{i=1}^n \left(\sum_{j=k}^m \langle \omega_i, \beta_j \rangle u_{j-1}\beta_j \right) \mu_i}.$$

Example 4.4. Let $M = SL_3(\mathbb{C})/B$ be the manifold of complete flags in $\mathbb{C}^3$, let $u = s_1 = 213$ and $v = s_1s_2s_1 = 321$. Then $\ell(321) = 3$, and

$$\Lambda_v^- = \alpha_1\alpha_2(\alpha_1 + \alpha_2).$$

There are two ascending chains of maximal length from u to v ,

$$\begin{aligned}\gamma_1 : \quad & s_1 \xrightarrow{*s_{\alpha_1+\alpha_2}} s_2 s_1 \xrightarrow{*s_{\alpha_2}} s_1 s_2 s_1 \\ \gamma_2 : \quad & s_1 \xrightarrow{*s_{\alpha_2}} s_1 s_2 \xrightarrow{*s_{\alpha_1}} s_1 s_2 s_1 ,\end{aligned}$$

and the contributions of the two chain are

$$\begin{aligned}E_\mu(\gamma_1) &= \alpha_1 \alpha_2 (\alpha_1 + \alpha_2) \cdot \frac{\mu_1 + \mu_2}{\alpha_2 \mu_1 + (\alpha_1 + \alpha_2) \mu_2} \cdot \frac{\mu_2}{\alpha_1 \mu_2} = \frac{\alpha_2 (\alpha_1 + \alpha_2) (\mu_1 + \mu_2)}{\alpha_2 \mu_1 + (\alpha_1 + \alpha_2) \mu_2} \\ E_\mu(\gamma_2) &= \alpha_1 \alpha_2 (\alpha_1 + \alpha_2) \cdot \frac{\mu_2}{\alpha_2 \mu_1 + (\alpha_1 + \alpha_2) \mu_2} \cdot \frac{\mu_1}{\alpha_2 \mu_1} = \frac{\alpha_1 (\alpha_1 + \alpha_2) \mu_2}{\alpha_2 \mu_1 + (\alpha_1 + \alpha_2) \mu_2} .\end{aligned}$$

Then

$$\tau_u(v) = E_\mu(\gamma_1) + E_\mu(\gamma_2) = \alpha_1 + \alpha_2 .$$

5. LIMITS AND CHAIN CONTRIBUTIONS

Note that, in Example 4.4, both $E_\mu(\gamma_1)$ and $E_\mu(\gamma_2)$ are rational expressions in α 's and μ 's, but their sum is a polynomial expression in the α variables only. In this section we show that we can eliminate the μ variables in (8) by sending them to 0, one component at a time.

The contribution (5) of the chain (4) can be written as

$$E_\mu(\gamma) = \Lambda_v^- \prod_{k=1}^m \frac{Q_\mu(u_{k-1}, u_k, v)}{\alpha(u_{k-1}, u_k)} ,$$

where, for $p \prec q \preceq r$ in W , we define

$$(9) \quad Q_\mu(p, q, r) = \frac{p\eta - q\eta}{p\eta - r\eta} = \frac{\sum_{i=1}^n (p\omega_i - q\omega_i)\mu_i}{\sum_{i=1}^n (p\omega_i - r\omega_i)\mu_i} .$$

Lemma 5.1. For $p, q \in W$, let

$$h(p, q) = \min\{i \mid p\omega_i \neq q\omega_i\} ,$$

with the convention that $h(p, p) = \infty$.

(1) If $p^{-1}q = s_{i_1}s_{i_2} \cdots s_{i_k}$ is a reduced decomposition, then

$$h(p, q) = \min\{i_1, \dots, i_k\} .$$

(2) If $p \prec q \prec r$ and $p\omega_i = r\omega_i$, then $p\omega_i = q\omega_i = r\omega_i$.

(3) If $p \prec q \prec r$, then

$$h(p, r) \leq h(p, q) \quad \text{and} \quad h(q, r) \leq h(p, q) .$$

(4) If $q = ps_\beta$ with $\beta \succ 0$ and $p\beta \succ 0$, then $h(p, q) = h(\beta)$, where, for nonzero $\beta = \beta^1\alpha_1 + \cdots + \beta^n\alpha_n$ in $\mathfrak{t}^*$, we define $h(\beta) = \min\{i \mid \beta^i \neq 0\}$.

Proof. Let $j = h(p, q)$ and ω_i a fundamental weight. Then (7) becomes

$$(10) \quad \omega_i - p^{-1}q\omega_i = \delta_{i_1, i} \alpha_{i_1} + \delta_{i_2, i} s_{i_1} \alpha_{i_2} + \cdots + \delta_{i_k, i} s_{i_1} s_{i_2} \cdots s_{i_{k-1}} \alpha_{i_k}.$$

If $i < j$, then $\omega_i = p^{-1}q\omega_i$, and since all the nonzero terms in (10) are positive roots, it follows that s_i can not appear in the reduced decomposition of $p^{-1}q$. If $i = j$, then $\omega_j \neq p^{-1}q\omega_j$, hence s_j must appear in the decomposition of $p^{-1}q$. Therefore $j = h(p, q)$ is the minimal letter that appears in a reduced word for $p^{-1}q$ and $p^{-1}q$ is in the subgroup of W generated by $s_j, s_{j+1}, \dots, s_n$.

For every $i = 1, \dots, n$ we have $p\omega_i - r\omega_i = (p\omega_i - q\omega_i) + (q\omega_i - r\omega_i)$, and since $p \prec q \prec r$, both $p\omega_i - q\omega_i$ and $q\omega_i - r\omega_i$ are in the non-negative cone generated by the simple roots. Therefore, if $p\omega_i - r\omega_i = 0$, then $p\omega_i - q\omega_i = 0$ and $q\omega_i - r\omega_i = 0$. Then $\{i \mid p\omega_i = r\omega_i\} \subset \{i \mid p\omega_i = q\omega_i\}$, which implies $h(p, r) \leq h(p, q)$. Similarly we get $h(p, r) \leq h(q, r)$.

If $q = ps_\beta$, then

$$\{i \mid p\omega_i \neq ps_\beta\omega_i\} = \{i \mid \omega_i \neq s_\beta\omega_i\} = \{i \mid (\omega_i, \beta) \neq 0\} = \{i \mid \beta^i \neq 0\},$$

hence $h(p, q) = h(\beta)$. $\square$

Consequently, if the coefficient of some μ_i in the denominator is zero, then the coefficient of μ_i in the numerator is also zero. This implies that

$$Q_\mu^0(p, q, r) \stackrel{\text{def}}{=} \lim_{\mu_1 \rightarrow 0} \left(\lim_{\mu_2 \rightarrow 0} \left(\cdots \left(\lim_{\mu_n \rightarrow 0} Q_\mu(p, q, r) \right) \cdots \right) \right)$$

is well defined. Moreover,

$$Q_\mu^0(p, q, r) = \begin{cases} \frac{p\omega_i - q\omega_i}{p\omega_i - r\omega_i}, & \text{if } h(p, r) = h(p, q) = i \\ 0, & \text{if } h(p, r) < h(p, q). \end{cases}$$

In particular, let $p = u_{k-1}$, $q = u_k = u_{k-1}\beta_k$, and $r = v$. Then

$$\frac{Q_\mu^0(u_{k-1}, u_k, v)}{\alpha(u_{k-1}, u_k)} = \begin{cases} \frac{\langle \omega_i, \beta_k \rangle}{u_{k-1}\omega_i - v\omega_i}, & \text{if } h(u_{k-1}, v) = h(u_{k-1}, u_k) = i \\ 0, & \text{if } h(u_{k-1}, v) < h(u_{k-1}, u_k). \end{cases}$$

Therefore

$$E(\gamma) \stackrel{\text{def}}{=} \lim_{\mu_1 \rightarrow 0} \left(\lim_{\mu_2 \rightarrow 0} \left(\cdots \left(\lim_{\mu_n \rightarrow 0} E_\mu(\gamma) \right) \cdots \right) \right)$$

is well-defined, and is zero whenever $h(u_{k-1}, v) < h(u_{k-1}, u_k)$ for some k . If $h(u_{k-1}, u_k) = h(u_{k-1}, v) = i_k$ for all $k = 1, \dots, m$, then

$$E(\gamma) = \Lambda_v^- \prod_{k=1}^m \frac{\langle \omega_{i_k}, \beta_k \rangle}{u_{k-1}\omega_{i_k} - v\omega_{i_k}} \neq 0.$$

We now take a closer look at the set of chains

$$\mathcal{C}_0(u, v) = \{\gamma \in \Sigma(u, v) \mid h(u_{k-1}, u_k) = h(u_{k-1}, v) \text{ for all } k = 1, \dots, m\}$$

with non-zero contribution after taking the limits. These chains have a simple description.

Lemma 5.2. Let $\gamma \in \Sigma(u, v)$ be the ascending chain of maximal length

$$\gamma : u = u_0 \xrightarrow{*s\beta_1} u_1 \xrightarrow{*s\beta_2} \dots \xrightarrow{*s\beta_m} u_m = v .$$

Then $\gamma \in \mathcal{C}_0(u, v)$ if and only if $h(\beta_1) \leq h(\beta_2) \leq \dots \leq h(\beta_m)$.

Proof. If $\gamma \in \mathcal{C}_0(u, v)$, then, using Lemma 5.1, we get

$$h(\beta_{k+1}) = h(u_k, u_{k+1}) = h(u_k, v) \geq h(u_{k-1}, v) = h(u_{k-1}, u_k) = h(\beta_k)$$

for all $k = 1, \dots, m-1$, proving one implication.

Conversely, let $\gamma \in \Sigma(u, v)$ such that $h(\beta_1) \leq h(\beta_2) \leq \dots \leq h(\beta_m)$. Let $k \in [m]$ be fixed. If $i < h(\beta_k)$, then $i < h(\beta_j) = h(u_{j-1}, u_j)$ for all $j \geq k$, hence $u_{j-1}\omega_i = u_j\omega_i$ for all $j \geq k$. Therefore $u_{k-1}\omega_i = u_m\omega_i = v\omega_i$, hence $h(u_{k-1}, v) \geq h(\beta_k) = h(u_{k-1}, u_k)$. But $h(u_{k-1}, v) \leq h(u_{k-1}, u_k)$ by Lemma 5.1, and the double inequality implies that $h(u_{k-1}, v) = h(u_{k-1}, u_k)$ for all $k \in [m]$, and therefore $\gamma \in \mathcal{C}_0(u, v)$. $\square$

Example. Returning to Example 4.4, we see that

$$E(\gamma_1) = \lim_{\mu_1 \rightarrow 0} \left(\lim_{\mu_2 \rightarrow 0} \frac{\alpha_2(\alpha_1 + \alpha_2)(\mu_1 + \mu_2)}{\alpha_2\mu_1 + (\alpha_1 + \alpha_2)\mu_2} \right) = \alpha_1 + \alpha_2 \neq 0$$

$$E(\gamma_2) = \lim_{\mu_1 \rightarrow 0} \left(\lim_{\mu_2 \rightarrow 0} \frac{\alpha_1(\alpha_1 + \alpha_2)\mu_2}{\alpha_2\mu_1 + (\alpha_1 + \alpha_2)\mu_2} \right) = 0 ,$$

as expected, since, for γ_1 we have $h(\alpha_1 + \alpha_2) = 1 \leq 2 = h(\alpha_2)$ and for γ_2 we have $h(\alpha_2) = 2 \not\leq 1 = h(\alpha_1)$.

6. POSITIVE FORMULA USING CHAINS

Up to this point, there have been no restrictions regarding the order in which the simple roots $\alpha_1, \dots, \alpha_n$ are listed. The next results, however, requires that the simple roots be listed in a specific order, as shown in the following Dynkin diagrams:

$$(11) \quad \begin{array}{ll} \text{Type A:} & \alpha_1 \text{ --- } \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \text{ --- } \alpha_n \\ \text{Type B:} & \alpha_1 \text{ --- } \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \rightrightarrows \alpha_n \\ \text{Type C:} & \alpha_1 \text{ --- } \alpha_2 \text{ --- } \dots \text{ --- } \alpha_{n-1} \leftrightsquigarrow \alpha_n \end{array}$$

We are now ready to formulate the main result of this paper. Let $\mathcal{C}_0(u, v)$ be the set of ascending chains of maximal length

$$(12) \quad \gamma : u = u_0 \xrightarrow{*s\beta_1} u_1 \xrightarrow{*s\beta_2} u_2 \rightarrow \dots \rightarrow u_{m-1} \xrightarrow{*s\beta_m} u_m = v$$

that satisfy the condition

$$(13) \quad h(\beta_1) \leq h(\beta_2) \leq \dots \leq h(\beta_m) ,$$

and for each such a chain γ , let

$$(14) \quad E(\gamma) = \Lambda_v^- \prod_{k=1}^m \frac{\langle \omega_{i_k}, \beta_k \rangle}{u_{k-1}\omega_{i_k} - v\omega_{i_k}},$$

where $i_k = h(\beta_k)$ for $k = 1, \dots, m$.

Theorem 6.1. If $u, v \in W$ then:

[Part I] The restriction of τ_u at v is given by

$$(15) \quad \tau_u(v) = \sum_{\gamma \in \mathcal{C}_0(u, v)} E(\gamma) = \sum_{\gamma \in \mathcal{C}_0(u, v)} \Lambda_v^- \prod_{k=1}^m \frac{\langle \omega_{i_k}, \beta_k \rangle}{u_{k-1}\omega_{i_k} - v\omega_{i_k}}.$$

[Part II] If the simple roots are ordered as in (11), then:

In types A and C:

$$E(\gamma) \in \mathbb{Z}_{\geq 0}[\alpha_1, \dots, \alpha_n],$$

hence (15) is a positive integral formula for computing $\tau_u(v)$.

In type B:

$$E(\gamma) \in \frac{1}{2^m} \mathbb{Z}_{\geq 0}[\alpha_1, \dots, \alpha_n],$$

hence (15) is a positive, but only rational, formula for $\tau_u(v)$.

[Part III] If the simple roots are ordered as in (11), then:

In type A: For every u and v , there is a particular choice of a reduced word I for v , and a bijection $F_I: \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ such that $E(\gamma) = SC(F_I(\gamma), I)$. In other words, the positive formula using chains is an alternative formulation of Billey's positive formula using subwords.

In type C: There exist u and v such that, whichever reduced word I for v we choose, the number of subwords in $\mathcal{R}(u, I)$ is not the same as the number of chains in $\mathcal{C}_0(u, v)$. Therefore, in type C, the two positive formulas are not manifestly equivalent.

Proof. The right hand side of the Goldin-Tolman formula (3) is independent of the particular choice of η in the positive chamber (and hence of μ with strictly positive components). Each term in the right hand side is well-defined after sending the components of μ to zero, one at a time in reverse order. If $\gamma \notin \mathcal{C}_0(u, v)$, then the limit is zero, and if $\gamma \in \mathcal{C}_0(u, v)$, then the limit is $E(\gamma)$, and this proves the statement in Part I.

With the ordering (11), if β is a positive root and $h(\beta) = i$, then $\langle \omega_i, \beta \rangle$ is a positive integer, so the product in the numerator of (14) is a positive integer. To prove the statements of Part II, we show that:

- (1) Each factor in the denominator of (14) cancels (over $\mathbb{Z}_{>0}$ in types A and C, and over $\frac{1}{2}\mathbb{Z}_{>0}$ in type B) one of the factors in Λ_v^- , and
- (2) The factors in the denominator of (14) are mutually independent.

These statements are consequences of the following technical lemma, which can be verified through straightforward computations.

Lemma 6.2. If $u \in W$ and $j = h(id, u)$, then:

In type A: u has a reduced decomposition of the form

$$(16) \quad u = s_k s_{k-1} \cdots s_j w$$

with $j \leq k \leq n$ and $h(id, w) > j$.

In type B or C: u has either a reduced decomposition (16) or a reduced decomposition of the form

$$(17) \quad u = s_t s_{t+1} \cdots s_{n-1} s_n s_{n-1} \cdots s_{j+1} s_j w ,$$

with $j \leq t < n$ and $h(id, w) > j$, but not both.

If u has a decomposition of the form (16), then

$$\omega_j - u\omega_j = \begin{cases} \alpha_j + \cdots + \alpha_k, & \text{in types A, C for } k \leq n \\ & \text{and type B if } k < n \\ \alpha_j + \cdots + \alpha_{n-1} + 2\alpha_n, & \text{in type B for } k = n \end{cases}$$

If u has a decomposition of the form (17) with $t > j$, then

$$\omega_j - u\omega_j = \begin{cases} \alpha_j + \cdots + \alpha_{t-1} + 2(\alpha_t + \cdots + \alpha_n), & \text{in type B} \\ \alpha_j + \cdots + \alpha_{t-1} + 2(\alpha_t + \cdots + \alpha_{n-1}) + \alpha_n, & \text{in type C} \end{cases}$$

If u has a decomposition of the form (17) with $t = j$, then

$$\omega_j - u\omega_j = \begin{cases} 2(\alpha_j + \cdots + \alpha_n) & , \text{ in type B} \\ 2(\alpha_j + \cdots + \alpha_{n-1}) + \alpha_n & , \text{ in type C} \end{cases}$$

If u is not of the form $s_j \cdots s_n \cdots s_j w$ in type B, then $\omega_j - u\omega_j$ is a root. In the excepted case, $\omega_j - u\omega_j$ is twice a root. In all cases $h(\omega_j - u\omega_j) = j$, and no distinct factors are multiples of each other.

The above lemma implies that

$$u_{k-1}\omega_{i_k} - v\omega_{i_k} = u_{k-1}(\omega_{i_k} - u_{k-1}^{-1}v\omega_{i_k})$$

is either a root (in types A and C) or, possibly, twice a root (in type B). From Lemma 4.3 we conclude that each factor of (14) cancels (over $\mathbb{Z}_{>0}$ in types A and C, and over $\frac{1}{2}\mathbb{Z}_{>0}$ in type B) one of the factors in Λ_v^- .

To finish the proof of Part II, we need to show that the factors are mutually independent. Suppose that is not true, and that for some chain $\gamma \in \mathcal{C}_0(u, v)$ and some $k < t$, the vectors $u_{k-1}\omega_{i_k} - v\omega_{i_k}$ and $u_{t-1}\omega_{i_t} - v\omega_{i_t}$ are dependent. Then $\omega_{i_k} - v^{-1}u_{k-1}\omega_{i_k}$ and $\omega_{i_t} - v^{-1}u_{t-1}\omega_{i_t}$ are also dependent. Using Lemma 6.2 we get that

$$i_k = h(\omega_{i_k} - v^{-1}u_{k-1}\omega_{i_k}) = h(\omega_{i_t} - v^{-1}u_{t-1}\omega_{i_t}) = i_t ,$$

hence $i_k = i_t$. Let $j = i_k = i_t$ be the common value. Since $\omega_j - v^{-1}u_{k-1}\omega_j$ and $\omega_j - v^{-1}u_{t-1}\omega_j$ are dependent, Lemma 6.2 implies that they must be equal, and therefore $u_{k-1}\omega_j - u_{t-1}\omega_j = 0$. Then the proof of Lemma 4.3 implies that $\langle \omega_j, \beta_k \rangle = 0$, and that is a contradiction, since $j = i_k = h(\beta_k)$. Therefore the factors in the denominator of (14) are independent, and that finishes the proof of Part II.

If $u = s_2$ and $v = s_1s_2s_1$ in B_2 , then the chain

$$\gamma : s_2 \xrightarrow{*s_{\alpha_1+2\alpha_2}} s_1s_2 \xrightarrow{*s_{\alpha_1+\alpha_2}} s_1s_2s_1$$

is in $\mathcal{C}_0(u, v)$, but

$$E(\gamma) = \alpha_1(\alpha_1 + \alpha_2)(\alpha_1 + 2\alpha_2) \cdot \frac{1}{2(\alpha_1 + \alpha_2)} \cdot \frac{1}{\alpha_1 + 2\alpha_2} = \frac{1}{2} \alpha_1,$$

hence in type B the formula (15) is not necessarily integral, and the denominators can be higher powers of 2.

The following example proves the Part III statement in type C .

Let $u = s_1$ and $v = s_1s_2s_1s_2$ in C_2 . Then $I_1 = [1, 2, 1, 2]$ and $I_2 = [2, 1, 2, 1]$ are the only reduced words for v , and each of them has two subwords that are reduced words for u . Hence whether $I = I_1$ or $I = I_2$, there are two subwords in $\mathcal{R}(u, I)$, hence two terms in (2), the positive formula using subwords.

There are four ascending chains of maximal length from u to v :

$$(18) \quad \begin{aligned} \gamma_1 : & s_1 \xrightarrow{*s_{\alpha_1+\alpha_2}} s_2s_1 \xrightarrow{*s_{\alpha_1+2\alpha_2}} s_1s_2s_1 \xrightarrow{*s_{\alpha_2}} s_1s_2s_1s_2 \\ \gamma_2 : & s_1 \xrightarrow{*s_{\alpha_1+\alpha_2}} s_2s_1 \xrightarrow{*s_{\alpha_2}} s_2s_1s_2 \xrightarrow{*s_{\alpha_1}} s_1s_2s_1s_2 \\ \gamma_3 : & s_1 \xrightarrow{*s_{\alpha_2}} s_1s_2 \xrightarrow{*s_{\alpha_1+\alpha_2}} s_2s_1s_2 \xrightarrow{*s_{\alpha_1}} s_1s_2s_1s_2 \\ \gamma_4 : & s_1 \xrightarrow{*s_{\alpha_2}} s_1s_2 \xrightarrow{*s_{\alpha_1}} s_1s_2s_1 \xrightarrow{*s_{\alpha_2}} s_1s_2s_1s_2 \end{aligned}$$

but only γ_1 satisfies (13), hence $\mathcal{C}_0(u, v) = \{\gamma_1\}$ has only one chain.

All that remains to be proved is the Part III statement in type A . In Section 7 we give an explicit version of the positive formula (23) in type A , in Section 8 we construct a general map connecting chains and subwords, and we finish the proof in Section 9. $\square$

The main result of Lemma 6.2 is that if $h(id, u) = j$, then $\omega_j - u\omega_j$ is a root (in type A and C) or a multiple of a root (in type B , and, as one can check, for G_2). However, in type D or for F_4 , $\omega_j - u\omega_j$ is a sum of positive roots that is not a multiple of a root, and therefore, the corresponding chain contribution (14) is positive but not even polynomial.

7. TYPE A POSITIVE FORMULA USING CHAINS

We apply the general result of Theorem 6.1 to $M = SL_n(\mathbb{C})/B$.

If $h(\beta) = k$, then $\beta = \alpha_{kj}$ for some $j \geq k$ and $\langle \omega_k, \beta \rangle = 1$. Hence all the factors in the numerator of (14) are equal to 1. The denominators can be computed using the following lemma.

Lemma 7.1. Let $u \in S_n$. If $h(id, u) = j$, then

$$(19) \quad \omega_j - u\omega_j = x_j - x_{u(j)} .$$

Proof. Let $u = s_k s_{k-1} \cdots s_j w$ be the decomposition (16). Then

$$\omega_j - u\omega_j = \alpha_j + \cdots \alpha_k = x_j - x_{k+1} ,$$

and since $h(id, w) > j$, we have $k+1 = u(j)$. $\square$

If γ is the ascending chain of maximal length (12) as in Theorem 6.1, then

$$u_{k-1}\omega_{i_k} - v\omega_{i_k} = u_{k-1}(\omega_{i_k} - u_{k-1}^{-1}v\omega_{i_k}) = x_{u_{k-1}(i_k)} - x_{v(i_k)} ,$$

and therefore

$$E(\gamma) = \left[\prod_{(a,b) \in \text{Inv}(v)} (x_a - x_b) \right] \prod_{k=1}^m \frac{1}{x_{u_{k-1}(i_k)} - x_{v(i_k)}} .$$

We summarize the results of this section and formulate our version of a type A positive formula for $\tau_u(v)$. Let $C_0(u, v)$ be the set of ascending chains of maximal length

$$(20) \quad \gamma: u = u_0 \xrightarrow{*(i_1, j_1)} u_1 \xrightarrow{*(i_2, j_2)} u_2 \rightarrow \cdots \rightarrow u_{m-1} \xrightarrow{*(i_m, j_m)} u_m = v$$

that satisfy the condition

$$(21) \quad i_1 \leq i_2 \leq \cdots \leq i_m ,$$

where $*(i, j)$ means that we are multiplying to the right by the transposition that swaps i and j . For such a chain γ , let

$$\text{Inv}(v, \gamma) = \text{Inv}(v) \setminus \{(u_{k-1}(i_k), v(i_k)) \mid k = 1, \dots, m\}$$

and

$$(22) \quad E(\gamma) = \prod_{(a,b) \in \text{Inv}(v, \gamma)} (x_a - x_b) .$$

Theorem 7.2. The restriction of τ_u at v is given by

$$(23) \quad \tau_u(v) = \sum_{\gamma \in C_0(u, v)} E(\gamma) = \sum_{\gamma \in C_0(u, v)} \prod_{(a,b) \in \text{Inv}(v, \gamma)} (x_a - x_b) .$$

Example 7.3. Consider the permutations $u = 2143$ and $v = 3421$ in S_4 . There are two chains in $\mathcal{C}_0(2143, 3421)$:

$$\begin{aligned}\gamma_1 : \quad & 2143 \xrightarrow{*(1,4)} 3142 \xrightarrow{*(2,3)} 3412 \xrightarrow{*(3,4)} 3421 \\ \gamma_2 : \quad & 2143 \xrightarrow{*(1,4)} 3142 \xrightarrow{*(2,4)} 3241 \xrightarrow{*(2,3)} 3421 .\end{aligned}$$

For the first chain $u_0 = 2143$, $u_1 = 3142$, $u_2 = 3412$, and $u_3 = v = 3421$. The edge $2143 \xrightarrow{*(1,4)} 3142$ deletes the factor $x_{u_0(1)} - x_{v(1)} = x_2 - x_3$. The edge $3142 \xrightarrow{*(2,3)} 3412$ deletes the factor $x_{u_1(2)} - x_{v(2)} = x_1 - x_4$. The edge $3412 \xrightarrow{*(3,4)} 3421$ deletes the factor $x_{u_2(3)} - x_{v(3)} = x_1 - x_2$. Therefore

$$\begin{aligned}E(\gamma_1) &= \frac{(x_2 - x_3)(x_1 - x_3)(x_2 - x_4)(x_1 - x_4)(x_1 - x_2)}{(x_2 - x_3)(x_1 - x_4)(x_1 - x_2)} = \\ &= (x_1 - x_3)(x_2 - x_4) = (\alpha_1 + \alpha_2)(\alpha_2 + \alpha_3) .\end{aligned}$$

Similarly

$$\begin{aligned}E(\gamma_2) &= \frac{(x_2 - x_3)(x_1 - x_3)(x_2 - x_4)(x_1 - x_4)(x_1 - x_2)}{(x_2 - x_3)(x_1 - x_4)(x_2 - x_4)} = \\ &= (x_1 - x_3)(x_1 - x_2) = (\alpha_1 + \alpha_2)\alpha_1 .\end{aligned}$$

Hence

$$\tau_{2143}(3421) = E(\gamma_1) + E(\gamma_2) = (\alpha_1 + \alpha_2)(\alpha_1 + \alpha_2 + \alpha_3) .$$

8. CHAINS AND SUBWORDS

In this section we give a general construction connecting ascending chains and subwords. This construction is valid for all Weyl groups and orderings of simple roots.

Let $u \prec v$ and let $I = [i_1, \dots, i_k]$ be a reduced word for v . Let $\mathcal{A}(u, v)$ be the set of all ascending chains

$$(24) \quad \gamma: u = u_0 \xrightarrow{*s_{\beta_1}} u_1 \xrightarrow{*s_{\beta_2}} u_2 \rightarrow \dots \rightarrow u_{m-1} \xrightarrow{*s_{\beta_m}} u_m = v ,$$

from u to v , not necessarily of maximal length, and let $\mathcal{C}(u, v)$ be the subset of $\mathcal{A}(u, v)$ consisting of the chains that satisfy $h(\beta_1) \leq \dots \leq h(\beta_m)$. Then

$$\mathcal{C}(u, v) \cap \Sigma(u, v) = \mathcal{C}_0(u, v) \subset \mathcal{C}(u, v) \subset \mathcal{A}(u, v) .$$

Let $\mathcal{S}(u, I)$ be the set of subwords of I that generate words for u by deleting the zeroes; then $\mathcal{R}(u, I)$ is a subset of $\mathcal{S}(u, v)$. We define a function $F_I: \mathcal{A}(u, v) \rightarrow \mathcal{S}(u, I)$ as follows.

Let γ be the ascending chain (24). We use the edges of γ , in reverse order, to delete letters from I and get a subword of I that is a word for u . Since

$u_{m-1} = vs_{\beta_m} \prec v$, the Strong Exchange Property implies that there exists a j (unique, since I is reduced) such that

$$u_{m-1} = s_{i_1} \cdots s_{i_{j-1}} s_{i_{j+1}} \cdots s_{i_k},$$

hence $I_{m-1} = [i_1, \dots, i_{j-1}, 0, i_{j+1}, \dots, i_k] \in \mathcal{S}(u_{m-1}, I)$ is a subword of I that is a word for u_{m-1} . Similarly, $u_{m-2} = u_{m-1}s_{\beta_{m-1}}$ and $\ell(u_{m-2}) < \ell(u_{m-1})$, so there exists a subword of I_{m-1} , obtained by deleting exactly one letter from I_{m-1} , that is a word for u_{m-2} . However, since $\ell(u_{m-1})$ is not necessarily equal to $\ell(v) - 1$, the word I_{m-1} may be non-reduced, and in this case the uniqueness of the deleted letter is not guaranteed. We choose the subword for which the deleted letter is the rightmost choice, and get a subword I_{m-2} of I_{m-1} (hence of I), that is a word, not necessarily reduced, for u_{m-2} . Continuing this process, with the same rule for making a choice, if needed, we get a sequence $I_{m-1}, \dots, I_0$ of subwords of I such that $I_k \in \mathcal{S}(u_k, I)$ and I_k is obtained from I_{k+1} by deleting one letter. We define $F_I(\gamma) = I_0$, the last subword in the sequence.

Example 8.1. Let $u = s_1$ and $v = s_1 s_2 s_1 s_2$ in the Weyl group of the root system C_2 . Let $I = [1, 2, 1, 2]$ be a reduced word for v and let

$$\gamma_1 : s_1 \xrightarrow{*s_{\alpha_1+\alpha_2}} s_2 s_1 \xrightarrow{*s_{\alpha_1+2\alpha_2}} s_1 s_2 s_1 \xrightarrow{*s_{\alpha_2}} s_1 s_2 s_1 s_2,$$

be an ascending chain of maximal length from u to v . Since $s_{\alpha_1+\alpha_2} = s_1 s_2 s_1$ and $s_{\alpha_1+2\alpha_2} = s_2 s_1 s_2$, the subword $F_I(\gamma_1)$ is computed as follows:

$$\begin{aligned} [1, 2, 1, 2] \cdot [2] &= [1, 2, 1, 0] = I_2 \\ [1, 2, 1, 0] \cdot [2, 1, 2] &= [0, 2, 1, 0] = I_1 \\ [0, 2, 1, 0] \cdot [1, 2, 1] &= [0, 0, 1, 0] = I_0 \end{aligned}$$

hence $F_I(\gamma_1) = [0, 0, 1, 0]$. The chains in $\Sigma(u, v)$ are given in (18), and $F_I(\gamma_1) = F_I(\gamma_2) = F_I(\gamma_3) = [0, 0, 1, 0]$ and $F_I(\gamma_4) = [1, 0, 0, 0]$.

Note that if $\gamma \in \Sigma(u, v)$, then $\ell(u_{k-1}) = \ell(u_k) - 1$ for all k , hence the words $I_{m-1}, \dots, I_1, I_0$ are reduced. Therefore at each stage there is only one possibility for the deleted letter and no choice is necessary.

Lemma 8.2. Let $\gamma \in \mathcal{A}(u, v)$ be an ascending chain. Then

$$F_I(\gamma) \in \mathcal{R}(u, I) \iff \gamma \in \Sigma(u, v).$$

Moreover, if $F_I : \mathcal{C}(u, v) \rightarrow \mathcal{S}(u, I)$ is bijective, then $F_I : \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ is also bijective.

Proof. Let $\gamma \in \mathcal{A}(u, v)$. Then $F_I(\gamma) \in \mathcal{R}(u, I)$ if and only if $F_I(\gamma)$ has $\ell(u)$ non-zero letters. This happens if and only if we delete $\ell(v) - \ell(u)$ letters from I , and since every deleted letter corresponds to an edge of γ , if and only if γ has $\ell(v) - \ell(u)$ edges, in other words, if and only if γ is an ascending chain

of maximal length. If $\gamma \in \mathcal{C}_0(u, v)$ then $\gamma \in \Sigma(u, v)$, hence $F_I(\gamma) \in \mathcal{R}(u, I)$. Therefore $F_I: \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ is well-defined.

Suppose that $F_I: \mathcal{C}(u, v) \rightarrow \mathcal{S}(u, I)$ is bijective. Then the restriction of F_I to $\mathcal{C}_0(u, v)$ is injective. Moreover, if $J \in \mathcal{R}(u, I)$, then there exists $\gamma \in \mathcal{C}(u, v)$ such that $F_I(\gamma) = J$. But then $\gamma \in \mathcal{C}(u, v) \cap \Sigma(u, v) = \mathcal{C}_0(u, v)$, hence $F_I: \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ is also surjective. $\square$

9. EQUIVALENCE OF POSITIVE FORMULAS

We show that in the case $G = SL_n(\mathbb{C})$, the positive formula (23) is an alternative formulation of Billey's formula (2), by proving that, for a particular choice of a reduced word I for v , the function F_I restricts to a bijection $F_I: \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ and $SC(F_I(\gamma), I) = E(\gamma)$ for all $\gamma \in \mathcal{C}_0(u, v)$.

Let $I(v)$ be the reduced word for v constructed inductively using (16). Then

$$I(v) = [I_1, I_2, \dots, I_{n-1}] ,$$

such that, for all $j = 1, \dots, n-1$,

- $I_j = [k_j, k_j - 1, \dots, j + 1, j]$ or $I_j = []$, and
- $h(s_{I_1} \cdots s_{I_j}, v) = h(id, s_{I_j} \cdots s_{I_1} v) = h(id, s_{I_{j+1}} \cdots s_{I_{n-1}}) > j$.

Example 9.1. If $v = s_2 s_1 s_3 s_2 s_3$, then $I_1 = [2, 1]$, $I_2 = [3, 2]$ and $I_3 = [3]$.

We can now prove the equivalence of our positive formula (using chains) with Billey's positive formula (using subwords).

Theorem 9.2. Let $u, v \in S_n$ and let $I = I(v)$. Then $F_I: \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ is a bijection and

$$SC(F_I(\gamma), I) = E(\gamma)$$

for all chains $\gamma \in \mathcal{C}_0(u, v)$.

Proof. In [Za] we proved that $F_I: \mathcal{C}(u, v) \rightarrow \mathcal{S}(u, I)$ is a bijection. Then, by Lemma 8.2, the map $F_I: \mathcal{C}_0(u, v) \rightarrow \mathcal{R}(u, I)$ is also a bijection.

To prove that $SC(F_I(\gamma), I) = E(\gamma)$, we show that the factor of Λ_v^- canceled by an edge of γ is the same as the factor canceled by the corresponding deleted letter from I .

In [Za] we also proved that if

$$(25) \quad \gamma: u = u_0 \xrightarrow{*s_{\beta_1}} u_1 \xrightarrow{*s_{\beta_2}} u_2 \rightarrow \cdots \rightarrow u_{m-1} \xrightarrow{*s_{\beta_m}} u_m = v ,$$

is a chain in $\mathcal{C}_0(u, v)$, then:

- (1) The edge $u_{k-1} \xrightarrow{*s_{\beta_k}} u_k$ deletes a letter from $I_{h(\beta_k)}$, and
- (2) The letters deleted in I_j are deleted from left to right.

Let $u_{k-1} \rightarrow u_k = u_{k-1}s_{\beta_k}$ be an edge of γ and let $j = h(\beta_k)$.

Suppose that $h(\beta_k) = \dots = h(\beta_{k+q-1}) = j < h(\beta_{k+q})$, hence β_k is the q^{th} occurrence of j from right to left. Let

$$J_k = [I_1, \dots, I_{j-1}, I'_j, I'_{j+1}, \dots, I'_{n-1}] \in \mathcal{R}(u_k, I(v))$$

be the subword obtained after using $s_{\beta_m}, \dots, s_{\beta_{k+1}}$ to delete letters from $I(v)$, and let

$$J_{k-1} = [I_1, \dots, I_{j-1}, I''_j, I'_{j+1}, \dots, I'_{n-1}] \in \mathcal{R}(u_{k-1}, I(v))$$

be the subword obtained after using s_{β_k} to delete one more letter from J_k . Then I'_j is of the form

$$I'_j = [k_j, \dots, \widehat{j_1}, \dots, \widehat{j_{q-1}}, \dots, j] \subset I_j,$$

and

$$I''_j = [k_j, \dots, \widehat{j_1}, \dots, \widehat{j_{q-1}}, \dots, \widehat{j_q}, \dots, j] \subset I'_j \subset I_j,$$

with $j_1 > j_2 > \dots > j_q$, since the letters in I_j are deleted from left to right.

The factor of Λ_v^- canceled by the edge $u_{k-1} \rightarrow u_k$ is

$$\begin{aligned} u_{k-1}\omega_j - v\omega_j &= (u_{k-1}\omega_j - u_k\omega_j) + \dots + (u_{k+q-2}\omega_j - u_{k+q-1}\omega_j) = \\ &= \langle \omega_j, \beta_k \rangle u_{k-1}\beta_k + \dots + \langle \omega_j, \beta_{k+q-1} \rangle u_{k+q-2}\beta_{k+q-1} = \\ &= s_{I_1} \dots s_{I_{j-1}} s_{k_j} \dots s_{j_1+1} (\widehat{s_{j_1}} \dots \widehat{s_{j_2}} \dots \widehat{s_{j_q}} \alpha_{j_q} + \dots + \alpha_{j_1}). \end{aligned}$$

But if $j_1 > j_2$, then

$$\widehat{s_{j_1}} \dots \widehat{s_{j_2}} \alpha_{j_2} + \alpha_{j_1} = s_{j_1} \dots \widehat{s_{j_2}} \alpha_{j_2},$$

and then, by induction on q ,

$$\widehat{s_{j_1}} \dots \widehat{s_{j_2}} \dots \widehat{s_{j_q}} \alpha_{j_q} + \dots + \alpha_{j_1} = s_{j_1} \dots s_{j_{q-1}} \dots \widehat{s_{j_q}} \alpha_{j_q}.$$

Therefore

$$u_{k-1}\omega_j - v\omega_j = s_{I_1} \dots s_{I_{j-1}} s_{k_j} \dots s_{j_q+1} \alpha_{j_q},$$

and that is precisely the factor of Λ_v^- canceled by the missing letter j_q , letter that has been deleted by the edge $u_{k-1} \rightarrow u_k$.

This finishes the proof of Theorem 9.2, which was the last piece in the proof of Theorem 6.1. We have therefore shown that for $G = SL_n(\mathbb{C})$, our positive formula (using chains) is an alternative version of Billey's positive formula (using subwords). $\square$

REFERENCES

- [BB] Björner, Anders and Francesco Brenti. *Combinatorics of Coxeter groups*. Graduate Texts in Mathematics, 231. Springer, New York, 2005.
- [Bi] Billey, Sara C. Kostant polynomials and the cohomology ring for G/B . *Duke Math. J.* **96** (1999), no. 1, 205–224. MR1663931 (2000a:14060)
- [GT] Goldin, Rebecca and Sue Tolman. A Generalized Schubert Calculus. Preprint. To appear in *Journal of Symplectic Geometry*.
- [Kn] Knutson, Allen. *A Littelmann-type formula for Duistermaat-Heckman measures*. Invent. Math., vol. **135** No. 1, 1999.
- [ST] Sabatini, Silvia and Sue Tolman. A Simpler Path Formula for Equivariant Cohomology. Preprint.
- [Za] Zara, Catalin. Chains, subwords, and fillings: strong equivalence of three definitions of the Bruhat order. *Electron. J. Combin.* 13 (2006), no. 1, Note 5, 13 pp. (electronic). MR2212512 (2006k:05222)

DEPARTMENT OF MATHEMATICS, UMASS BOSTON, 100 MORRISSEY BLVD., BOSTON, MA 02125